Three-Dimensional Astronomical Tomography and Wavefront Decomposition: Physical Evaluation of Circumstellar Shell Inversion, Stokes Polarimetry, and AROS Telemetry Pipelines

Target Identification and Morphological Dichotomy

The operational directive specifies target architectures consisting of concentric circumstellar shells and irregular dwarf galaxy morphological structures, referencing the Galaxies Losing Oxygen via Winds (GLOW) survey and the irregular dwarf galaxy NGC 4449. However, empirical examination of the telemetry frame reveals a fundamental astrophysical discrepancy: the astronomical image corresponds directly to the Egg Nebula (CRL 2688 / AFGL 2688), a prototypical carbon-rich protoplanetary nebula situated in the constellation Cygnus at a heliocentric distance of approximately 1.0\text{ kpc} (3.1 \times 10^{16}\text{ km}).

The visual morphology displayed in the image is defined by four prominent, collimated radial illumination beams ("searchlight beams") that cross a series of concentric arcs and limb-brightened ripples centered on an obscured asymptotic giant branch (AGB) progenitor. These optical beams originate where starlight escapes through bipolar cavities excavated in a dense, equatorial dust torus, illuminating outer concentric dust shells that were episodically shed during thermal pulsations of the central supergiant.

The caption embedded within the incoming telemetry conflates this circumstellar architecture with archival data from NGC 4449, a Magellanic-type irregular starburst dwarf galaxy situated in Canes Venatici at a distance of 3.82 \pm 0.27\text{ Mpc}. The GLOW investigation, utilizing Hubble Space Telescope (HST) instrumentation, is designed to quantify galactic-scale chemical feedback and baryonic cycles. Specifically, GLOW measures how galactic winds generated by massive young star clusters and supernova blast waves expel synthesized elements into the circumgalactic medium, demonstrating that low-mass dwarf galaxies retain on average only 11\% of their produced oxygen.

A rigorous computational pipeline requires an explicit separation of the physical regimes governing these disparate astronomical structures. Circumstellar shells in protoplanetary nebulae represent coherent mass-ejection fronts driven by a localized central stellar engine, regulated by dust nucleation kinetics and homologous expansion across physical scales of 10^{-2}\text{ pc} to 10^{-1}\text{ pc}. In contrast, dwarf irregular starbursts encompass stochastic, multi-source star-forming complexes spanning several kiloparsecs, where diffuse gas depletion is mediated by super-bubbles expanding at velocities exceeding 200\text{ km s}^{-1} into the intergalactic environment.

Physical Parameter	Circumstellar Architecture (Egg Nebula / CRL 2688)	Extragalactic Starburst Architecture (NGC 4449 / GLOW)
Object Classification	Bipolar Protoplanetary Nebula (PPN)	Magellanic Dwarf Irregular Galaxy (dIrr)
Heliocentric Distance (D)	\sim 1.0\text{ kpc} (\approx 3,260\text{ ly})	3.82 \pm 0.27\text{ Mpc} (\approx 12.5\text{ Mly})
Angular Scale Conversion	1.0" \approx 1,000\text{ AU} (1.5 \times 10^{11}\text{ km})	1.0" \approx 18.5\text{ pc} (5.7 \times 10^{14}\text{ km})
Total Physical Span	\sim 0.3\text{ pc} (6 \times 10^4\text{ AU})	\sim 6.2\text{ kpc} across optical disk
Central Driving Source	Carbon-rich post-AGB star (F5\text{Ia}) + binary companion	Distributed massive young star clusters (10^3 - 10^5\text{ M}_\odot)
Dominant Emission Mechanism	Dust scattering of central starlight (P_{\text{lin}} \le 70\%)	Direct stellar continuum and collisionally excited line emission
Kinematic Velocity Scale	v_{\text{exp}} \sim 15 - 30\text{ km s}^{-1} (halo); \le 600\text{ km s}^{-1} (jets)	Super-bubble expansion v \sim 220\text{ km s}^{-1}; galactic winds
Primary Inversion Vector	Polarimetric Stokes and Doppler morpho-kinematics	Stellar population synthesis and synthetic CMD modeling

Radiative Transfer Inversion and Line-of-Sight Depth Stratification

Optical Depth Formulation and Stokes Parameter Inversion

Reconstructing an unprojected three-dimensional volume density distribution \rho(x, y, z) from a two-dimensional projected optical intensity map I_{\text{obs}}(x, y) requires resolving the monochromatic radiative transfer equation along the line-of-sight Z-axis:

$$I_\nu(x, y) = \int_{-\infty}^{\infty} S_\nu(x, y, z) \, e^{-\tau_\nu(x, y, z)} \, \kappa_\nu \rho(x, y, z) \, dz$$

In this formulation, S_\nu represents the local source function, \kappa_\nu denotes the mass extinction opacity, and \tau_\nu(x, y, z) = \int_{z}^{\infty} \kappa_\nu \rho(x, y, z') \, dz' expresses the optical depth through the intervening medium toward the observer.

Within a cold circumstellar reflection envelope such as CRL 2688, where in-situ thermal dust emission is negligible across optical and near-infrared bandpasses (\lambda \le 2\ \mu\text{m}), the source function is governed by the scattering of central stellar flux F_*(\nu) by sub-micron carbonaceous grains:

$$S_\nu(x, y, z) = \frac{\omega_\nu F_*(\nu)}{4\pi \, r(x, y, z)^2} \, \Phi(\theta_{\text{sca}}) \, e^{-\tau_*(\mathbf{r})}$$

where \omega_\nu represents the single-scattering albedo, r = \sqrt{x^2 + y^2 + z^2} is the radial distance from the central stellar core, \tau_*(\mathbf{r}) is the radial attenuation from the star to the scattering element, and \Phi(\theta_{\text{sca}}) is the scattering phase function parameterized by the Henyey-Greenstein asymmetry factor g = \langle \cos \theta_{\text{sca}} \rangle.

Linear polarimetry executed with the Advanced Camera for Surveys (ACS) or Wide Field Camera 3 (WFC3) breaks the spatial depth degeneracy along the line of sight. When unpolarized stellar continuum photons undergo single scattering off dust grains, the scattered radiation becomes linearly polarized perpendicular to the scattering plane. The Stokes vector \mathbf{S} = [I, Q, U, V]^T quantifies this polarization signature:
P_{\text{lin}}(x, y) = \frac{\sqrt{Q(x, y)^2 + U(x, y)^2}}{I(x, y)} \theta_{\text{pol}}(x, y) = \frac{1}{2} \arctan \left(\frac{U(x, y)}{Q(x, y)} \right)
For sub-micron dust distributions where grain radii satisfy the condition a \ll \lambda, the fractional polarization for single-scattered light exhibits an intense dependence on the scattering angle \theta_{\text{sca}}, reaching a sharp maximum at \theta_{\text{sca}} = 90^\circ:
P(\theta_{\text{sca}}) = P_{\text{max}} \frac{\sin^2 \theta_{\text{sca}}}{1 + \cos^2 \theta_{\text{sca}}}
The physical geometry connecting the projected radial coordinate R_{\text{proj}} = \sqrt{x^2 + y^2}, the line-of-sight coordinate z, and the scattering angle is constrained by \cos \theta_{\text{sca}} = z / \sqrt{R_{\text{proj}}^2 + z^2}. Inverting the observed degree of linear polarization determines the line-of-sight depth vector z for discrete, optically thin shell features:
\vert{}z(x, y)\vert{} = R_{\text{proj}} \left[\frac{1 - \sqrt{1 - \left(\frac{P_{\text{lin}}(x, y)}{P_{\text{max}}} \right)^2}}{\frac{P_{\text{lin}}(x, y)}{P_{\text{max}}}} \right]^{1/2}
High-resolution imaging polarimetry of CRL 2688 records polarization levels between 50\% and 70\% throughout both lobes. This provides direct physical verification that scattering occurs near \theta_{\text{sca}} \approx 90^\circ \pm 20^\circ, establishing that the primary bipolar reflection lobes reside nearly in the plane of the sky (z \approx 0) with an inclination angle of 10^\circ to 20^\circ.

Kinematic Spectroscopic Inversion

Complementing polarimetric depth inversion, high-dispersion long-slit and grism spectroscopy conducted with the Space Telescope Imaging Spectrograph (STIS) and Cosmic Origins Spectrograph (COS) resolves the velocity distribution v_{\text{los}}(x, y) along the line of sight. By targeting diagnostic emission transitions—such as shock-excited molecular hydrogen (H_2\ v=1\text{--}0\ S(1) at 2.1218\ \mu\text{m}) or optical forbidden lines ([O I] \lambda 6300, [N II] \lambda 6584, [S II] \lambda\lambda 6716, 6731)—the recorded Doppler shift reflects the projected line-of-sight component of nebular expansion:
v_{\text{los}}(x, y, z) = v_r(r) \frac{z}{r} + v_{\text{sys}}
In circumstellar envelopes undergoing radial, ballistic outflow, the expansion velocity follows a linear Hubble-type relation (v_r[span_108](start_span)[span_108](end_span) \propto r). Under this condition, Doppler shifts map directly to line-of-sight depth:
z = \left(\frac{v_{\text{los}} - v_{\text{sys}}}{v_{\text{exp}}} \right) R_{\text{max}}
This spatio-kinematic mapping separates overlapping forward and rear shell boundaries, decoupling approaching and receding outflow fronts and resolving front-to-back density contrasts across the envelope.

Temporal Metric Equivalence and Light-Travel Geometry

Treating spatial intensity morphology as a phase-encoded temporal metric requires evaluating two distinct physical mechanisms operating across disparate timescales:
The dynamic mass-loss timescale correlates the physical separation of concentric arcs \Delta R directly with the temporal recurrence period of circumstellar mass ejections \Delta

t_{pulse} via the radial wind velocity $v_{\[span_112](start_span)[span_112](end_span)[span_113](start_span)[span_113](end_span)text{wind}}$:

$$\Delta t_{\text{pulse}} = \frac{\Delta R}{v_{\text{wind}}} = \frac{D \cdot \Delta \theta}{v_{\text{wind}}}$$

In the Egg Nebula, observational measurements reveal an average angular arc spacing of $\Delta \theta \approx 0.3''\te[span_114](start_span)[span_114](end_span)[span_116](start_span)[span_116](end_span)[span_118](start_span)[span_118](end_span)xt{--}0.7''$. At an established distance of $D = 1.0\text{ kpc}$, this angular separation equates to a physical shell interval of $\Delta R \approx (4.5\text{--}10.5) \times 10^{15}\text{ cm}$ $(300\text{--}700[span_115](start_span)[span_115](end_span)[span_117](start_span)[span_117](end_span)[span_119](start_span)[span_119](end_span)\text{ AU})$. Combining this distance with millimeter CO rotational line measurements that indicate an AGB wind velocity of $v_{\text{wind}} \approx 15\text{--}19\text{ km s}^{-1}$, the interval between successive mass-loss pulses is calculated as:

$$\Delta t_{\text{pulse}} \approx \frac{(4.5\text{--}10.5) \times 10^{10}\text{ km}}{17\text{ km s}^{-1}} \approx 84\text{ to }196\text{ years}$$

This calculated periodicity matches the characteristic thermal pulse timescales (helium shell flashes) predicted by stellar evolution models for intermediate-mass stars at the tip of the asymptotic giant branch, as well as periodic orbital perturbations induced by a close binary companion.

On prompt timescales, light-travel time effects across circumstellar dust structures introduce geometric arrival delays. Photons emitted by the central star that scatter toward the observer obey the geometric relation defining a paraboloid of equal arrival time:

$$c \, t = r - z = \sqrt{R_{\text{proj}}^2 + z^2} - z$$

Solving this relation for physical depth z establishes that scattered wavefront arrivals from the far side of the nebula ($z < 0$) are delayed relative to forward-scattered photons ($z > 0$):

$$z(R_{\text{proj}}, t) = \frac{R_{\text{proj}}^2 - (ct)^2}{2ct}$$

Consequently, transient luminosity variations of the central engine create expanding scattering echoes across the concentric shells, encoding the three-dimensional geometry directly into the time-variable surface brightness profiles.

Algorithmic Processing and Processing Directives

The operational execution pipeline comprises four integrated processing stages that transform raw telemetry into calibrated three-dimensional data products:

1. **PSF Deconvolution and Artifact Mitigation:** Raw telemetry frames from HST WFC3, ACS, and STIS undergo calibration, dark-frame subtraction, and flat-field division, followed by iterative Richardson-Lucy restoration against synthesized point-spread function libraries.

2. **Spatial and Kinematic Inversion:** Deconvolved intensity fields are processed in parallel through Stokes polarimetric inversion (mapping linear polarization fractions to scattering angles and line-of-sight depths), Doppler spectroscopic slicing (extracting radial velocity matrices from slit and grism spectra), and multi-epoch astrometric alignment (measuring arc expansion velocities).
3. **Harmonic Radial Decomposition:** Polar coordinate transformations of the restored spatial intensity fields isolate azimuthally integrated radial profiles, which are detrended and processed via Fast Fourier Transforms to quantify structural periodicity.
4. **Volumetric Data-Cube Assembly:** Inverted spatial, kinematic, and polarimetric coordinates are mapped into a standardized three-dimensional Cartesian manifold \Lambda(x, y, z), parameterized by localized velocity vectors and optical depth vectors.

PSF Deconvolution and Bayesian Inverse Modeling

Hubble optical imaging of circumstellar nebulae is limited by diffraction artifacts, wide-angle scattered light halos, and detector cross-talk, which obscure low-surface-brightness arcs located near bright central illumination cones. To resolve inner concentric shells without introducing high-frequency noise amplification, an accelerated Richardson-Lucy iterative deconvolution is applied:

$$\hat{I}^{(n+1)}(x, y) = \hat{I}^{(n)}(x, y) \left[\frac{I_{\text{obs}}(x, y)}{\hat{I}^{(n)}(x, y) \otimes \text{PSF}(x, y)} \otimes \text{PSF}^T(x, y) \right]$$

The point spread function \text{PSF}(x, y) is generated from optical modeling libraries (such as STScI *Tiny Tim*) or empirical point-source calibrations matching the specific detector filters and focal plane coordinates. To suppress edge ringing and noise divergence after numerous iterations, the pipeline integrates a Bayesian maximum a posteriori (MAP) framework with Total Variation (TV) regularization:

$$\hat{I}_{\text{MAP}} = \arg \max_{I \ge 0} \left\{ \ln \mathcal{L}(I_{\text{obs}} \vert{} I) - \lambda_{\text{reg}} \int \int \vert{}\nabla I(x, y)\vert{} \, dx \, dy \right\}$$

This regularization suppresses noise in smooth circumstellar zones while preserving sharp edge discontinuities across expanding dust fronts.

Radial Frequency Decomposition via Fast Fourier Transforms

To quantitatively test whether circumstellar shell ejection is modulated by underlying periodic resonance phenomena, the deprojected two-dimensional intensity map I(x, y) is transformed into polar coordinates I(R_{\text{proj}}, \phi) centered on the stellar coordinates. Azimuthal integration over angular sectors exhibiting clean shell visibility yields a one-dimensional radial surface brightness profile \bar{I}(r):

$$\bar{I}(r) = \frac{1}{\phi_2 - \phi_1} \int_{\phi_1}^{\phi_2} I(r, \phi) \, d\phi$$

Geometric flux dilution is removed by subtracting a baseline power-law trend I_{\text{trend}}(r) \propto r^{-2}, isolating radial density perturbations \delta I(r) = \bar{I}(r) - I_{\text{trend}}(r). The spatial frequency spectrum is evaluated by applying a one-dimensional Fast Fourier Transform (FFT) combined with a Tukey windowing function W(r) to mitigate boundary truncation leakage:

$$\mathcal{P}(k_r) = \left\vert{} \int_{r_{\text{in}}}^{r_{\text{out}}} \delta I(r) \, W(r) \, e^{-i 2\pi k_r r} \, dr \right\vert{}^2$$

Statistically significant spectral peaks isolated at a fundamental spatial frequency k_0 and associated integer harmonics (2k_0, 3k_0) confirm coherent, clock-like episodic ejections. A broadband power spectrum devoid of discrete peaks indicates stochastic hydrodynamical

fragmentation or turbulent boundary layer shredding rather than periodic mass-loss episodes.

Morpho-Kinematic Reconstruction and Predictive Forward Modeling

In standard astrophysical modeling, deterministic three-dimensional reconstruction is conducted using morpho-kinematic frameworks such as SHAPE (Steffen et al.). These tools synthesize Doppler velocity profiles with spatial intensity contours to generate three-dimensional polygonal or analytical surface meshes governed by physical mass conservation:
$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{v}) = 0$$
Forward-looking kinetic modeling over secular baselines of $+10^3$ to $+10^4\text{ years}$ incorporates ballistic momentum conservation modulated by ram-pressure deceleration against the ambient interstellar medium:
$$\mathbf{r}(t + \Delta t) = \mathbf{r}(t) + \mathbf{v}(\mathbf{r}) \Delta t - \frac{1}{2} \left(\frac{\rho_{\text{ISM}} v^2}{\rho_{\text{shell}} \Delta r_{\text{shell}}} \right) \hat{\mathbf{v}} \, (\Delta t)^2$$
Where neural architectures are integrated, they operate as Physics-Informed Neural Networks (PINNs) that embed the Navier-Stokes hydrodynamic equations within the network loss functions, guaranteeing that predicted shell dispersion adheres strictly to energy and momentum conservation.

Technical Audit of AROS Provenance and Protocol Enhancements

Infrastructure and Operating Provenance

The operational directive specifies deployment within the AROS™ infrastructure (Albert and Robot's Operating System, accessible via aros.albertlane.net and dev.albertlane.net), governed by proprietary software licensing constraints.
Technical evaluation of the hosting topography reveals that dev.albertlane.net operates as an immutable, content-addressed snapshot archive where prior build deployments are permanently version-locked. The internal service mesh comprises containerized Docker microservices running Python FastAPI, Spring, and NGINX reverse proxies within a local network (box/local), routed to external endpoints via Cloudflare Workers. Cryptographic integrity is maintained via AES-GCM encrypted local storage keys and SHA-256 build verification hashes, ensuring tamper-evident tracking chains for processed scientific data.
Incoming data packets from edge nodes pass through Cloudflare Workers routing gates into the local service mesh. Inside the local environment, the NGINX gateway handles reverse proxy routing, directing compute tasks to FastAPI execution microservices. The execution engines read and write to AES-GCM encrypted local key-value storage, after which completed calculations are compiled into immutable, content-addressed snapshots signed with SHA-256 Merkle hashes.

Physical Evaluation of AROS Protocol Enhancements

The specialized analytical directives introduced within the AROS protocol can be evaluated against established observational astrophysics and instrumentation standards:

Protocol 1 (Dynamic Noise-Floor Suppression via Stochastic Modeling) is technically valid and operationally necessary. Optical charge-coupled devices on HST (ACS/WFC and WFC3/UVIS) are susceptible to radiation-induced charge transfer inefficiency (CTI) trails, cosmic ray residues, and read-noise fluctuations in faint outer fields. Implementing stochastic background suppression—utilizing Generalized Extreme Value (GEV) noise modeling or Anscombe-transformed wavelet thresholding—effectively isolates real circumstellar arcs from spatial Poisson noise without attenuating faint structural flux.

Vector D ("Quantum State Interferometry") represents a nomenclature discrepancy relative to standard astrophysical practices. In stellar observational physics, macroscopic gas shells are not probed via quantum wave-state interference. However, the underlying physical objective—mapping localized thermal micro-fluctuations and thermodynamic ages along the line of sight—is routinely accomplished via **Multi-Transition Spectral Energy Distribution (SED) Inversion and Non-LTE Molecular Diagnostics**. By evaluating rotational-vibrational excitation states of molecular hydrogen (such as the ratio between the H_2\ v=1\text{--}0\ S(1) and v=2\text{--}1\ S(1) transitions) or high-J carbon monoxide transitions (^{12}\text{CO} / ^{13}\text{CO} ladders), local kinetic temperatures T_{\text{kin}} and excitation conditions can be mapped directly to distinct mass-loss epochs.

The proposed "Metric Wavefront Simulation" similarly represents a conceptual misnomer when applied to circumstellar scales. In weak-field Newtonian regimes surrounding intermediate-mass stars, general relativistic metric expansion perturbations (\Phi_{\text{grav}} / c^2 \sim G M_* / (r c^2) \le 10^{-8}) are completely negligible. The physical phenomenon that alters photon arrival wavefront geometries across line-of-sight depth zones is the **geometric light-travel delay across parabolic scattering surfaces** (light echoes) coupled with interstellar phase-screen scattering. Treating these wavefront interactions with classical geometric optics satisfies all empirical observational constraints without invoking non-existent spatial metric expansions.

Archival Data Synthesis and Instrumentation Matrix

Executing three-dimensional morpho-kinematic reconstruction across both circumstellar (Egg Nebula) and extragalactic (NGC 4449) architectures requires integrating complementary multi-instrument datasets from the Hubble Space Telescope archive.

Target System	HST Instrument	Operating Configuration	Primary Physical Observable	Pipeline Analytical Function
CRL 2688	**WFC3 / UVIS & IR**	F606W, F673N, F814W, F110W, F160W	High-contrast scattered starlight, concentric arcs, limb brightening	High-resolution morphology; radial FFT spatial frequency profiling
CRL 2688	*ACS / WFC*	POL0V, POL60V, POL120V + Broad Filters	Linear Stokes parameters (I, Q, U), degree of polarization P_{\text{lin}}	Inversion of single-scattering angles to compute line-of-sight depth z
CRL 2688	**STIS / FUV-MAMA**	Long-Slit Spectroscopy / Grism G230MB, G430M	Spatially resolved Doppler shifts, gas kinematics	Line-of-sight expansion velocity field v_{\text{los}}(z) calibration

Target System	HST Instrument	Operating Configuration	Primary Physical Observable	Pipeline Analytical Function
CRL 2688	**Multi-Epoch HST**	WFPC2 (1995) \rightarrow ACS (2002) \rightarrow WFC3 (2009/2026)	Proper motion displacement \mu = \D[span_195](start_span)[span_195](end_span)[span_196](start_span)[span_196](end_span)elta \theta / \Delta t of concentric arcs	Geometric expansion baseline (D = v_{\text{exp}} / \mu) calibration
NGC 4449	**ACS / WFC**	F435W (B), F555W (V), F814W (I[span_90](start_span)[span_90](end_span)[span_94](start_span)[span_94](end_span)[span_98](start_span)[span_98](end_span)), F658N (H\alpha)	Starburst cluster resolution, ionized gas filaments, super-bubbles	Resolved stellar population photometry and synthetic CMD fitting
NGC 4449	**COS / FUV**	G130M, G160M "down-the-barrel"	Interstellar absorption profiles, column densities N(\text{Si II}), N(\text{O I})	Metal-enriched galactic wind outflow velocities and mass-loss rates

Synthesis and Algorithmic Realization

The deconstruction of two-dimensional astronomical projections into calibrated three-dimensional depth vectors (Z-axis stratification) is an achievable and rigorous methodology when grounded in classical radiative transfer and spatio-kinematic principles. The target geometry supplied in the telemetry stream is conclusively identified as the protoplanetary Egg Nebula (CRL 2688), whose circumstellar architecture represents an ideal testbed for polarimetric Stokes inversion and radial Fourier harmonic decomposition.

Operational realization of the AROS Data Cube (\Lambda) requires synthesizing three independent coordinate axes:

1. Projected transverse sky coordinates (x, y) derived from sub-pixel, Richardson-Lucy deconvolved spatial intensity frames.
2. Depth coordinates (z) established via Stokes parameter inversion (Q, U \rightarrow \theta_{\text{sca}} \rightarrow z) across the reflection-dominated searchlight lobes.
3. Velocity coordinates (v_z) isolated via high-resolution slit spectroscopy (STIS/COS), mapping line-of-sight mass outflow and internal shock interfaces.

Translating theoretical pipeline enhancements into standard computational workflows requires defining "quantum state interferometry" as multi-transition non-LTE molecular line excitation

modeling, while replacing "metric wavefront simulations" with classical light-travel time delay paraboloid calculations.
When anchored to verifiable physical conservation laws, this analytical framework provides an end-to-end capability for reconstructing circumstellar mass-loss histories, validating thermal-pulse periodicity, and projecting morphological evolution across multi-millennial timescales.

Works cited

1. Stars Make Oxygen for Small Galaxies. They Lose Most of It., https://www.rutgers.edu/news/stars-make-oxygen-small-galaxies-they-lose-most-it 2. Hubble Reveals Cosmic Mystery, https://www.nasa.gov/image-article/hubble-reveals-cosmic-mystery/ 3. GLOW. I. Comprehensive Measurements of Gas-rich, Star-forming, https://www.researchgate.net/publication/417363240_GLOW_I_Comprehensive_Measurements_of_Gas-rich_Star-forming_Low-mass_Galaxies_in_the_Nearby_Universe 4. Near-infrared spectroscopy of (proto)-planetary nebulae: molecular, https://academic.oup.com/mnras/article/344/1/262/1010312 5. Proto-planetary Nebulae - Martin Schoenball, https://www.schoenball.de/astronomie/projekte/ppn/ppn_en.htm 6. Egg Nebula - NASA Science, https://science.nasa.gov/asset/hubble/egg-nebula/ 7. shows images of IRAS 171503224 derived from 40 and 480 s, https://www.researchgate.net/figure/shows-images-of-IRAS-171503224-derived-from-40-and-480-s-exposures-in-order-to_fig2_230918473 8. BIPOLAR AND MULTIPOLAR JETS IN PROTOPLANETARY AND, https://www.astroscu.unam.mx/rmaa/RMxAC..13/PDF/RMxAC..13_sahai.pdf 9. A Dust-Scattering Model for M1-92: A Revised Estimate of the Mass, https://www.mdpi.com/2075-4434/12/4/44 10. Stellar Fireworks Are Ablaze in Galaxy NGC 4449 - NASA Science, https://science.nasa.gov/asset/hubble/stellar-fireworks-are-ablaze-in-galaxy-ngc-4449/ 11. new HST/ACS observations of the irregular galaxy NGC 4449 - arXiv, https://arxiv.org/html/0708.0852v2 12. Hubble study finds dwarf galaxies retain little of their oxygen, https://www.spaceeve.com/blogs/space-news/hubble-glow-dwarf-galaxies-missing-oxygen 13. Nearby Dwarf Galaxies are Missing Most of Their Oxygen, Hubble Data Show, https://www.sci.news/astronomy/nearby-dwarf-galaxies-oxygen-hubble-15168.html 14. The birth of a planetary nebula around the carbon star IRC+10216, https://articles.adsabs.harvard.edu/pdf/1998MNRAS.300L..29S 15. 1984ApJ. . .278. .186Y The Astrophysical Journal, 278:186-194, https://adsabs.harvard.edu/pdf/1984ApJ...278..186Y 16. È2 km imaging polarimetry of CRL 2688 (the Egg Nebula) using (a, https://www.researchgate.net/figure/E2-km-imaging-polarimetry-of-CRL-2688-the-Egg-Nebula-using-a-NICMOS-Camera-2-after_fig1_2435578 17. arXiv:astro-ph/0601533v1 24 Jan 2006, https://arxiv.org/pdf/astro-ph/0601533 18. ST-ECF Newsletter - ESA/Hubble, https://esahubble.org/static/archives/stecfnewsletters/pdf/hst_stecf_0026.pdf 19. Summer School on Interstellar Processes: Abstracts of Contributed, https://ntrs.nasa.gov/api/citations/19870005688/downloads/19870005688.pdf?attachment=true 20. HST Cycle 33 Abstract Catalog - STScI, https://www.stsci.edu/files/live/sites/www/files/home/hst/proposing/approved-programs/_documents/HSTCycle33-Approved-Abstracts.pdf 21. Spatio-kinematics of the optical nebula M1-92 with HST/STIS,

https://www.researchgate.net/publication/262981045_Spatio-kinematics_of_the_optical_nebula_M1-92_with_HSTSTIS 22. Shape: A 3D Modeling Tool for Astrophysics - Semantic Scholar, https://www.semanticscholar.org/paper/Shape%3A-A-3D-Modeling-Tool-for-Astrophysics-Steffen-Koning/13e22f5362ec2df35b0b8cd7059ea3b6263e5131 23. A 3D Geometrical Model of Molecular Line Emission from Planetary, https://www.mdpi.com/2075-4434/14/4/65 24. 100-yr mass-loss modulations on the asymptotic giant branch, https://academic.oup.com/mnras/article/324/4/1117/1316774 25. II. The Carbon Star AFGL 3068 and the Planetary Nebulae NGC, https://academic.oup.com/mnras/article/495/2/2234/5838050 26. HubblesBeautifulUniverse 20250404 | PDF | Hubble Space Telescope, https://www.scribd.com/document/1068377410/HubblesBeautifulUniverse-20250404 27. Investigating the nature of the Fried Egg nebula: CO mm-line and, https://www.researchgate.net/publication/272381431_Investigating_the_nature_of_the_Fried_Egg_nebula_CO_mm-line_and_optical_spectroscopy_of_IRAS_17163-3907 28. 'Distance mapping' and the 3D structure of BD +30° 3639, https://academic.oup.com/mnras/article/423/1/925/1748275 29. SHAPE 2010, https://www.jb.man.ac.uk/meetings/APN5/pdfposters/32_steffen_poster.pdf 30. AROS immutable DEV, https://dev.albertlane.net/ 31. VEIL — albertlane.net, https://d1.albertlane.net/ 32. Albert Lane — Public Archive · SEC 17684-273-411-436, https://albertlane.net/ 33. ALMA Compact Array observations of the Fried Egg nebula - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC5300097/ 34. Molecular line emission in asymmetric envelopes of evolved stars, https://core.ac.uk/download/304640160.pdf