Empirical Architectures of Natural and Subterranean Intelligence: Decentralized Biology, Adaptive Cognition, and Survival Engineering

The phenomenon of intelligence on Earth is empirically characterized by thermodynamic efficiency, decentralized computation, and direct material adaptation. Across evolutionary timeframes, distributed biological networks, non-human organisms, and localized human societies have developed solutions to complex optimization problems without centralized computational architectures, bureaucratic hierarchies, or speculative capital incentives. These systems operate directly at the physical boundary layer, resolving challenges of resource titration, microclimate engineering, material manufacture, and navigation through recursive physical feedback loops grounded in evolutionary pressure, material physics, and empirical observation.

Decentralized Biological Problem-Solving and Dynamic Resource Allocation

Centralized processing architectures introduce latency, communication bottlenecks, and systemic vulnerability to localized damage. In contrast, distributed biological architectures execute complex decisions through local interfacial sensing, biochemical reaction-diffusion gradients, and morphological mechanics.

Subterranean Mycorrhizal Symbioses: Reciprocal Enforcement and Nutrient Flux Kinetics

Mycorrhizal symbioses, established over 400 to 450 million years of terrestrial plant evolution, constitute vast distributed biological trade networks that link vascular root systems across diverse ecosystems. These associations predominantly comprise arbuscular mycorrhizas (AM), which penetrate root cortical cells to form branched arbuscules, and ectomycorrhizas (EcM), which envelop root tips in an external fungal mantle and an intercellular Hartig net. The functional core of this mutualism involves the exchange of photosynthetically fixed host carbon—transported as hexoses and fatty acids—for fungal-acquired soil minerals, primarily orthophosphate (\text{P}_i) and inorganic nitrogen compounds.

Rather than operating as passive conduits that equalize concentration differences along indiscriminate source-sink gradients, these networks function through local, bidirectional reciprocal enforcement. Isotopic labeling and quantitative real-time PCR experiments utilizing *Medicago truncatula* and distinct arbuscular mycorrhizal fungal species (*Glomus intraradices* and *Glomus aggregatum*) have demonstrated that fungal symbionts selectively modulate nutrient delivery based on localized host carbon payments. Fungi absorb inorganic orthophosphate from the soil matrix beyond the root depletion zone—a boundary layer typically

extending 1 to 2 millimeters from the root surface—and condense the ions into long-chain polyphosphates (PolyP) within vacuoles to facilitate transport while preventing internal metabolic interference.

At the symbiotic periarbuscular interface, these chains are enzymatically cleaved into shorter polyphosphate fragments and exported across the fungal membrane into the interfacial apoplast only when the adjacent host plant cell provides reciprocal carbon rewards. Host plants dynamically alter their sucrose allocation, providing higher hexose efflux to hyphal branches that supply elevated nutrient volumes, while penalizing uncooperative or hoarding strains. When individual fungal branches experience carbon deprivation, they restrict phosphorus translocation to the non-cooperative host root, either redirecting it to alternative root nodes or storing it as long-chain PolyP. This decentralized economic titration maintains multi-species mutualisms across forest beds without centralized systemic regulation, stabilizing underground carbon drawdown and nutrient cycling across shifting climatic regimes.

Neuromuscular Autonomy and Peripheral Somatosensory Processing in Cephalopoda

The neuroanatomy of octopuses (*Octopus vulgaris*, *Octopus bimaculoides*) exhibits a marked divergence from the vertebrate paradigm of centralized cephalic governance. Out of approximately 5 \times 10^8 total neurons, roughly two-thirds are localized in the peripheral nervous system, primarily distributed along the axial nerve cords and brachial ganglia of the eight prehensile arms.

Anatomical Subsystem	Estimated Neuronal Proportion	Primary Functional Mandate
Central Brain (Supra- and Subesophageal Masses)	~35%–40%	High-level motor initiation, visual integration, spatial memory, affective state regulation
Brachial Ganglia & Axial Nerve Cords (Peripheral)	~60%–65%	Semi-autonomous motor coordination, local pattern generation, stereotypic reflex mediation
Sucker Rim Ganglionic Arrays	Dense sub-ganglionic clusters	Chemotactile sensory transduction, mechanosensory feedback, localized adhesion control

Each arm acts as a quasi-autonomous sensory-motor processing organ. Electrophysiological and behavioral assays show that an arm surgically or functionally isolated from the central supraesophageal brain preserves the capacity to perform complex motor programs. Such limbs can explore substrate fissures, execute searching trajectories, retrieve tactile objects, and withdraw from noxious chemical stimuli. This organization resolves the immense kinematic degrees of freedom inherent to a muscular hydrostat lacking rigid skeletal articulation.

Instead of the central brain computing individual trajectories for hundreds of independent suckers and complex muscle orientations—transverse, longitudinal, and helically arranged oblique bundles—central commands provide high-level recruitment signals. The peripheral brachial cord translates these commands locally using sensory feedback loops. Sucker epithelia possess specialized chemotactile receptors capable of detecting hydrophobic, non-volatile compounds and mechanical textures at the substrate interface, allowing decentralized parallel

processing across all eight arms simultaneously without central bottlenecking. This design demonstrates how complex physical problem-solving emerges from peripheral morphology coupled to localized neural networks.

Swarm Stigmergy and Thermodynamic Ventilation in Macrotermitinae Arcologies

The nest architecture of mound-building termites (*Macrotermes michaelseni*, *Odontotermes obesus*) provides an empirical model of decentralized thermodynamic regulation. Towering up to several meters above the ground surface, these structures house millions of workers and massive subterranean cultures of the symbiotic fungus *Termitomyces*. Maintaining internal homeostasis within these colonies requires continuous gas exchange to purge respiratory carbon dioxide and maintain relative humidity, all while stabilizing internal nest temperatures against external climatic fluctuations.

Earlier models proposed by Martin Lüscher posited that internal metabolic heat generated by the colony and fungus comb acted as a simple thermosiphon, lofting warm air upward through a central chimney and pulling it down through sealed outer conduits. However, in situ sensor measurements, tracer gas clearance assays, and thermal imaging conducted by Turner, Soar, and Mahadevan have overturned this paradigm. Termite mounds function not as steady heat engines, but as external respiratory systems that harvest ambient environmental oscillations. During daytime solar insolation, the mound's thin, highly exposed peripheral flutes heat rapidly due to low thermal inertia, while the voluminous subterranean core remains insulated and cool. This creates an upward convective flow in the outer flutes that draws cool, dense air down through the central chimney. At night, the process reverses: the flutes rapidly lose heat via radiative cooling to the sky, while the deep core retains heat from its thermal mass and metabolic activity. The resulting temperature inversion drives warm air upward through the central chimney, expelling accumulated carbon dioxide through the porous outer walls in a diurnal convective cycle.

The driving buoyancy pressure (ΔP) in these internal loops is governed by fluid thermal expansion:

$$\Delta P = \rho \alpha \Delta T g h$$

where ρ is air density, α is the thermal expansion coefficient of air ($\approx 1/T$), ΔT is the spatial temperature differential between peripheral flutes and the internal core, g is gravitational acceleration, and h is the vertical convective height. The resulting volumetric flow rate (Q) through the internal conduit networks is modeled by Poiseuille's law across open internal channels:

$$Q = \frac{\pi r^4 \Delta P}{8 \mu L}$$

where μ is dynamic air viscosity, r represents channel radius, and L is conduit length.

The construction of this thermodynamic organ occurs without executive direction. Termite workers operate via stigmergy, a decentralized coordination mechanism where individuals respond to local sensory cues—such as carbon dioxide concentration, air velocity, and moisture gradients—by depositing soil particles mixed with salivary secretions. By continually modifying the porosity and cross-sectional diameter of surface apertures in response to localized microclimatic stress, the collective adjusts its architectural boundary layer to maintain homeostatic equilibrium.

Non-Human Adaptive Cognition and Material

Traditions

Adaptive cognition in non-human animals demonstrates that foresight, causal comprehension, and culture-based transmission of material techniques are functional adaptations evolved to exploit complex ecological niches.

Avian Structural Fabrication and Causal Hierarchy in Corvids

Field and experimental ethology focusing on the New Caledonian crow (*Corvus moneduloides*) reveals sophisticated tool manufacturing and structural manipulation outside the mammalian lineage. These birds fabricate functional tools from living vegetation to extract wood-boring larvae from tree trunks.

Rather than relying on opportunistically collected debris, New Caledonian crows manufacture distinct tool morphotypes, including stepped implements cut from the barbed margins of *Pandanus* leaves (*Pandanus spiralis*, *Pandanus tectorius*) and hooked twigs harvested from dicotyledonous shrubs. During *Pandanus* tool fabrication, crows select live, turgid leaves, avoiding dried or decayed vegetation that splinter prematurely along structural fibers. The bird cuts into the leaf blade, tears longitudinally along the fiber lines, steps upward, and cuts again, producing tapered, multi-stepped implements with a wide grip base and a pointed, serrated working edge. Counterpart leaf cutouts documented across New Caledonia confirm that these geometric patterns are standardized and culturally conserved across regional populations.

When crafting hooked twig tools, crows snip off terminal nodes, strip adjacent foliage, and sculpt the remaining wood into an angled hook, adjusting the curvature to optimize mechanical extraction. In controlled cognitive testing, crows demonstrate hierarchical planning and physical reasoning. In multi-stage metatool assays, subjects use a short stick to extract an intermediate implement, which is then deployed to obtain a terminal food reward, organizing behaviors around mental models of means-end causality rather than conditioned motor reflexes. In water-displacement paradigms, crows drop dense, sinking stones into water columns to raise the water level to retrieve floating food, while ignoring hollow objects or sawdust substrates that do not follow liquid displacement physics.

Primate Lithic Archaeological Records: Diachronic Variation in Bearded Capuchins

Primate archaeological excavations at the Caju BPF2 site in Serra da Capivara National Park (SCNP), Brazil, have established that wild bearded capuchin monkeys (*Sapajus libidinosus*) maintain a persistent lithic tool tradition. Radiocarbon dating of stratified charcoal samples associated with buried stone artifacts demonstrates an unbroken sequence of lithic technology spanning between 2,400 and 3,000 calibrated years Before Present (\text{cal BP}).

Capuchins in this semi-arid Caatinga ecosystem deliberately select and transport heavy, hard quartzite cobbles from regional gravel beds to act as hammerstones, which they strike against flat sandstone boulders or horizontal tree root complexes that serve as anvils. This percussive technology is employed to crack hard, thick-shelled endemic food sources, including cashew nuts (*Anacardium occidentale*) and tough palm nuts (*Attalea barreirensis*, *Astrocaryum campestre*).

Chronological Phase	Calibrated Age Range	Primary Lithic Metrics & Utilized Material	Functional Resource Target
Phase IV (Basal Stratum)	~3,000 to 2,400 \text{cal BP}	High-mass quartzite cobbles; extensive battering scars and incipient percussion cones	Tough palm nuts, dense seeds, high-resistance tubers
Phase II / III (Middle)	~2,400 to 300 \text{cal BP}	Intermediate-mass hammerstones; concentrated anvil pitting and surface micro-flaking	Transitional foraging on mixed hard and moderately resistant flora
Phase I (Upper Stratum)	~100 \text{cal BP} to Present	Lower-mass, selected smooth quartzite pebbles; localized percussive crushing wear	Modern endemic cashew nuts (*Anacardium occidentale*)

The archaeological record demonstrates that this primate lithic industry has not remained static. Morphological analysis of artifacts across strata reveals systematic diachronic changes in tool selection: early capuchin populations utilized significantly larger, heavier hammerstones with severe percussion pitting, whereas modern populations utilize smaller cobbles adapted to the physical resistance of seasonal cashew nuts. The presence of uniform wear traces—such as point-impact striae, incipient cones of percussion, and crushed surfaces—distinguishes these tools from geologically fractured stones, documenting non-human technological adaptation across hundreds of generations.

Cetacean Epigenetic Material Culture: Shark Bay Dolphin Sponging Traditions

In Shark Bay, Western Australia, a subpopulation of Indo-Pacific bottlenose dolphins (*Tursiops aduncus*) exhibits specialized tool use termed "sponging". Individual dolphins detach marine basket sponges (*Dictyoceratida*) from the seafloor and position them over their rostra using their mouth. The conical sponge functions as a mechanical shield during benthic foraging in deep-water channels (>8\text{ meters} depth) composed of rough substrate and cherty gravel. The primary prey targeted via sponging are small, benthic barred sandperch (*Parapercis nebulosa*) that lack swim bladders. Because these teleosts have no air-filled bladders, they produce negligible acoustic echoes on dolphin echolocation arrays, forcing the animals to flush them manually from sediment beds. Foraging in these coarse sediments without a protective sponge cone causes significant tissue abrasion, lacerations, and exposure to toxic benthic fauna.

Long-term behavioral studies coupled with population genetics by Krützen et al. evaluated mitochondrial DNA control sequences and 12 biparentally inherited nuclear microsatellite markers across sponging and non-sponging dolphins.

The empirical findings confirm social rather than purely genetic or ecological transmission:

- Sponging behavior occurs within deep channels that are shared by non-sponging conspecifics, demonstrating that the behavior is not an obligatory response to ecological constraint.

- Pedigree and association data do not fit single-locus or multi-locus Mendelian inheritance models, ruling out a simple genetic basis.
- Sponging dolphins are unified by a single matrilineal mitochondrial haplotype (Haplotype H), demonstrating that this foraging specialization is transmitted culturally along maternal lines via social learning, where calves observe and practice the behavior over long developmental associations.

Species / Population	Functional Tool Type	Primary Mechanical Action	Information Transmission Mode	Diagnostic Physical Markers
New Caledonian Crow (*Corvus moneduloides*)	Stepped *Pandanus* leaves, hooked twigs	Tensile probing, crevice extraction, larval hooking	Individual trial with cumulative social facilitation	Standardized stepped margins on living vegetation
Bearded Capuchin (*Sapajus libidinosus*)	Quartzite hammerstones, sandstone anvils	High-impact vertical percussion, endocarp fracture	Observational social learning and cultural traditions	Percussive pits, micro-flake scars, anvil depressions
Bottlenose Dolphin (*Tursiops aduncus*)	Conical marine sponges (*Dictyoceratida*)	Benthic rostral shielding during gravel probing	Vertical matrilineal social transmission	Truncated sponge bodies, uniform rostral wear

Ground-Level Human Ingenuity: Survival-Driven Physics and Subterranean Material Science

Historic and indigenous technologies frequently solved acute environmental challenges through thermodynamic balancing, passive gravity management, and material biochemistry. These structures were developed out of survival necessity in marginal environments, providing sustainable infrastructure over centuries.

Subterranean Gravitational Hydraulics: The Civil Engineering of Qanat Systems

Originating on the arid Iranian plateau in the first millennium BCE, the *qanat* (alternatively known as *karez* or *foggara*) is an underground gravity-driven aqueduct engineered to transport water across long distances without mechanical pumping. The system taps groundwater in alluvial fans at the base of mountain ranges and conveys it through gently inclined subterranean tunnels to lowland agricultural and urban settlements.

Construction begins with the excavation of an exploratory "mother well" (*madar-chah*) into the upland alluvial water table, often reaching depths between 30 and 100 meters. Specialized well-diggers (*muqannis*) then excavate a subterranean gallery extending from the lowland daylight point toward the water source. Vertical shafts are sunk along the gallery axis every 20 to 50 meters to provide ventilation, allow removal of excavated debris, and maintain horizontal and vertical alignment.

The hydraulic gradient (S) of the subterranean tunnel is precisely controlled to balance fluid mechanics. The slope is maintained between 1:1{,}000 and 1:1{,}500 (0.05\%\text{ to }0.1\%). The mean flow velocity (v) of the water in these unlined alluvial channels is described by the Gauckler-Manning open-channel equation:

$$v = \frac{1}{n} R_h^{2/3} S^{1/2}$$
where:

- n represents the Manning roughness coefficient of the earthen or chiseled stone conduit ($\approx 0.025\text{–}0.035\text{ s}\cdot\text{m}^{-1/3}$),
- R_h is the hydraulic radius (A/P, cross-sectional flow area divided by wetted perimeter),
- S is the energy slope of the channel bed.

This gradient represents a critical hydraulic compromise:

- If the slope exceeds 1:1{,}000, flow velocity surpasses the critical threshold ($\approx 0.6\text{–}0.8\text{ m/s}$), creating hydrodynamic shear stress that erodes the unreinforced alluvial floor and causes the gallery to collapse.
- If the slope falls below 1:2{,}000, flow velocity drops below the transport velocity of suspended silt particles, leading to sedimentation, channel choking, and localized waterlogging.

By running water through deep subterranean conduits beneath an insulating mantle of dry soil, the qanat limits evaporation losses in arid regions where summer ambient temperatures exceed 40°C and potential evaporation exceeds $3{,}000\text{ mm/year}$. Powered solely by gravitational potential energy ($\Delta U = m g \Delta h$), the system provides baseflow without fossil energy inputs or mechanical maintenance. Because discharge depends on regional groundwater elevation, qanats dynamically adjust to climatic variations, preventing the rapid depletion that characterizes modern motorized tube-well extraction.

Microclimatic Energy Storage: The Waru-Waru Agroecosystems of the Lake Titicaca Basin

In the high-altitude Andean Altiplano ($3{,}800\text{ to }4{,}000\text{ meters}$ above sea level), pre-Columbian cultures (including the Pukara, Tiwanaku, and Colla) engineered extensive raised-field systems known as *waru-waru* (or *suka kollus*) across more than 82,000 hectares.
The Altiplano environment presents severe agricultural constraints: seasonal flooding during the summer rainy season, prolonged seasonal droughts, and radiation frosts during clear winter nights when ambient temperatures drop to between -5°C and -10°C.
The physical design of *waru-waru* consists of elevated agricultural berms (width 4–10 m, elevation 0.5–1.5 m) separated by wide, continuous canals (width 2–4 m, depth 0.5–1 m).
The system mitigates radiation frost through thermal storage and convective heat exchange:

- Liquid water has a high specific heat capacity: $c_w \approx 4.184 \times 10^3 \text{ J}\cdot\text{kg}^{-1}\cdot\text{K}^{-1}$ compared to dry organic soil, which has a specific heat capacity of: $c_s \approx 0.8\text{–}1.0 \times 10^3 \text{ J}\cdot\text{kg}^{-1}\cdot\text{K}^{-1}$.
- During intense daylight solar exposure, the canal water absorbs and stores thermal energy: $Q = m c_w \Delta T$
- At night, as cold air settles into low-lying areas, the warm canal water releases its stored energy to the air immediately above the beds through thermal convection, longwave radiation, and the generation of a light water-vapor layer.

Finite-element thermal modeling and field observations show this process creates a localized microclimatic thermal blanket, elevating canopy temperatures by 1°C to 3°C above ambient air, which prevents ice crystal formation within plant tissue.
The system also functions as a self-renewing nutrient cycle. The shallow canals cultivate aquatic vegetation, algae, and nitrogen-fixing cyanobacteria. Periodic dredging of this canal

sediment (*champa*) provides organic fertilizer that is reapplied to the adjacent planting beds, maintaining continuous soil productivity without fallow cycles or synthetic chemical inputs.

Civil Engineering System	Regional Context	Dominant Physical / Thermodynamic Mechanism	Key Structural Parameter
Qanat Subterranean Aqueducts	Iranian Plateau, Central Asian basins	Gravity-driven open channel flow below thermal boundary layer	Channel slope calibrated to 1:1{,}000\text{–}1:1{,}500; zero evaporation
Waru-Waru Raised Fields	Lake Titicaca Basin, Altiplano	Sensible heat absorption and nocturnal convective microclimate buffering	Canals 2–4 m wide; water maintains canopy 1^\circ\text{C}\text{–}3^\circ\text{C} above frost line
Terra Preta Anthrosols	Central Amazonian lowlands	Pyrogenic carbon matrix enhancing cation exchange capacity (CEC)	High biochar surface area; persistent functional group oxidation
Etak Oceanic Navigation	Polynesian / Micronesian open ocean	Celestial star-compass vectors and hydrodynamic swell diffraction	32-house sidereal azimuthal tracking without physical instrumentation

Pedological Pyrolysis and Oceanic Vector Wayfinding: Terra Preta and Siderial Navigation

Additional historic technologies highlight how localized material manipulation and cognitive mapping solved severe environmental challenges.

In the humid Amazon Basin, pre-Columbian agriculturalists transformed weathered, nutrient-poor soils (ferralsols and acrisols) into highly fertile *Terra Preta de Índio* (Amazonian dark earths). Under high tropical rainfall and temperatures, applied organic matter normally oxidizes rapidly and soil nutrients leach away. Indigenous land managers addressed this by incorporating fine-grained charcoal (biochar) produced through low-temperature, oxygen-depleted smoldering pyrolysis of organic waste, combined with animal bones and pottery fragments.

This pyrogenic carbon matrix provides an extensive internal surface area that resists microbial degradation. Over time, abiotic oxidation creates carboxyl and phenolic functional groups on the edges of the aromatic carbon sheets, increasing the soil's Cation Exchange Capacity (CEC). This matrix binds essential cations—including calcium, magnesium, potassium, and ammonium—preventing them from leaching during monsoonal rains. Modern soil analyses confirm that these anthrosols remain stable and fertile across centuries without chemical fertilization.

In maritime navigation, Polynesian and Micronesian wayfinders developed *etak*, a cognitive dead-reckoning framework that enabled open-ocean transits covering thousands of kilometers across the Pacific without instrumentation. The navigational paradigm integrates three physical and biological data vectors:

- **The Sidereal Star Compass (*Kāpehu Whetū*):** Navigators memorize the rising and setting points of specific stars along the horizon, establishing a directional compass

divided into 32 distinct azimuthal houses.
- **Hydrodynamic Swell Interference (*Keana Pua*):** Navigators sense the motion of the canoe hull relative to persistent oceanic swells generated by trade winds. By detecting swell refraction, deflection, and cross-swell interference patterns caused by distant islands, wayfinders determine the presence and direction of landmasses located beyond the visual horizon.
- **Biological Vector Transduction:** Navigators utilize the crepuscular foraging ranges of pelagic seabirds, such as the brown noddy (*Anous stolidus*) and white tern (*Gygis alba*). Because these birds fly directly out to sea in the morning and return to island roosts at dusk, their directional flight paths indicate landing trajectories within a 30 to 50 nautical mile radius.

Universal Axioms of Physical and Biological Computation

Analyzing decentralized biological structures, non-human material cultures, and indigenous engineering reveals fundamental principles that define real-world, functional intelligence:

- **Thermodynamic Coupling to Ambient Energy Fluxes:** Functional intelligence minimizes external energy consumption by coupling its structures directly to environmental gradients. Termite mounds leverage solar radiation cycles to drive convective ventilation; *waru-waru* fields exploit water's specific heat capacity to shield crops from radiation frosts; and *qanats* tap gravitational potential energy to transport groundwater across arid plains indefinitely. The intelligence of these systems resides in aligning architectural morphology with natural physical forces.
- **Localized Sensory-Motor Integration at the Boundary Layer:** Decentralized systems eliminate communication bottlenecks by placing sensory processing directly at the physical interface. In cephalopod arms, chemotactile sensing and neuromuscular responses are coordinated locally by the brachial ganglia; in mycorrhizal networks, nutrient-carbon trading rates are determined locally across cellular interfaces via dynamic biochemical feedback; and in New Caledonian crow tool crafting, birds make continuous modifications to leaf-strip curvature based on real-time feedback from plant fibers.
- **Long-Term Physical Fitness Over Institutional Scale:** Ground-level systems demonstrate their efficacy through material persistence under natural selection. Capuchin tool traditions have endured across 3,000 years of environmental change; *qanat* infrastructure has maintained continuous water flow across millennia; and *Terra Preta* soils continue to retain nutrients thousands of years after their formation.

These empirical systems demonstrate that intelligence is not defined by centralized control, administrative authority, or speculative models. Real-world intelligence is the ability of an organism, colony, or human community to observe ambient physical constraints, integrate sensory feedback at the boundary layer, and implement material solutions that maintain dynamic thermodynamic and ecological stability.

Works cited

1. (PDF) Mycorrhizal Networks - ResearchGate, https://www.researchgate.net/publication/417013402_Mycorrhizal_Networks 2. The Common Mycelial Network (CMN) of Forests,

https://clas.ucdenver.edu/ges/common-mycelial-network-cmn-forests 3. Common Mycorrhizae Network: A Review of the Theories and ... - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC10512311/ 4. Mechanisms of Arbuscular Mycorrhizal Mediation of Plant-Plant, https://beverlab.ku.edu/sites/beverlab/files/files/25.pdf 5. Reciprocal rewards stabilize cooperation in the mycorrhizal symbiosis, https://pubmed.ncbi.nlm.nih.gov/21836016/ 6. (PDF) Reciprocal Rewards Stabilize Cooperation in the Mycorrhizal, https://www.researchgate.net/publication/51566071_Reciprocal_Rewards_Stabilize_Cooperation_in_the_Mycorrhizal_Symbiosis 7. Soil to Cytoplasm: The Mycorrhizal Phosphorus Express and Its, https://www.mdpi.com/2309-608x/12/7/473 8. Mycorrhizal network - Wikipedia, https://en.wikipedia.org/wiki/Mycorrhizal_network 9. Octopus: Multiple minds or just a slow thinker?. Animal Sentience 26, https://www.wellbeingintlstudiesrepository.org/cgi/viewcontent.cgi?article=1513&context=animsent 10. Toward an Understanding of Octopus Arm Motor Control - PMC - NIH, https://pmc.ncbi.nlm.nih.gov/articles/PMC10755184/ 11. (PDF) Consciousness of octopuses—on their own terms, https://www.researchgate.net/publication/394897948_Consciousness_of_octopuses-on_their_own_terms 12. Solar-powered ventilation of African termite mounds - Soft Math Lab, https://softmath.seas.harvard.edu/publication/solar-powered-ventilation-of-african-termite-mounds/ 13. Solar-powered ventilation of African termite mounds, https://journals.biologists.com/jeb/article/220/18/3260/18709/Solar-powered-ventilation-of-African-termite 14. Climate control in termite mounds: A review and new insights from X, https://pure.hw.ac.uk/ws/portalfiles/portal/138451791/2406.13400v1.pdf 15. Can termites teach us to build environmentally friendly communities?, https://massivesci.com/articles/termite-mound-arcology-climate-control/ 16. (PDF) Beyond biomimicry: What termites can tell us about realizing, https://www.researchgate.net/publication/255650482_Beyond_biomimicry_What_termites_can_tell_us_about_realizing_the_living_building 17. How termites ventilate, https://seas.harvard.edu/news/how-termites-ventilate 18. Harvard study: termite mounds circulate air, sneezing once a day, https://www.harvardmagazine.com/2015/10/harvard-study-finds-that-termite-mound-functions-as-lung-that-sneezes-carbon-dioxide 19. The New Caledonian Crow Tool Use Observations/Experiments, https://english.arabpsychology.com/experiments/new-caledonian-crow-tool-use-gavin-hunt-russell-gray/ 20. Causal Reasoning in New Caledonian Crows | PDF | Force - Scribd, https://www.scribd.com/document/807039496/AB-C-2020-Vol7-3-Taylor 21. Gavin R. Hunt | University of Auckland | 56 Publications | 446 Citations, https://scispace.com/authors/gavin-r-hunt-53plizjsd1 22. [PDF] The development of pandanus tool manufacture in wild New, https://www.semanticscholar.org/paper/The-development-of-pandanus-tool-manufacture-in-New-Holzhaider-Hunt/f539e78be00700862db289fca808dd20a14fab11 23. Do new caledonian crows solve physical problems through causal, https://pubmed.ncbi.nlm.nih.gov/18796393/ 24. Complex cognition and behavioural innovation in New Caledonian, https://royalsocietypublishing.org/rspb/article/277/1694/2637/73251/Complex-cognition-and-behavioural-innovation-in 25. Three thousand years of wild capuchin stone tool use - ResearchGate, https://www.researchgate.net/publication/333982318_Three_thousand_years_of_wild_capuchin_stone_tool_use 26. Three thousand years of wild capuchin stone tool use - PubMed,

https://pubmed.ncbi.nlm.nih.gov/31235926/ 27. The Ergonomics of Stone Tool Use and Production - Apollo,
https://www.repository.cam.ac.uk/items/609ca5cf-0130-4332-bef6-0f44a2aaa54c 28. Serra da Capivara - NeoPReGo,
https://neoprego.org/research-sites-locais-de-pesquisa/serra-da-capivara/ 29. Stone tool use by wild capuchin monkeys (Sapajus libidinosus) at,
https://www.researchgate.net/publication/323831674_Stone_tool_use_by_wild_capuchin_monkeys_Sapajus_libidinosus_at_Serra_das_Confusoes_National_Park_Brazil 30. Cultural transmission of tool use in bottlenose dolphins - PNAS,
https://www.pnas.org/doi/10.1073/pnas.0500232102 31. Cultural transmission of animal tool use driven by trade-offs,
https://royalsocietypublishing.org/rsos/article/12/7/241900/235478/Cultural-transmission-of-animal-tool-use-driven-by 32. A culturally transmitted foraging technique in Indo-Pacific bottlenose,
https://royalsocietypublishing.org/rstb/article/381/1955/20250017/482594/A-culturally-transmitted-foraging-technique-in 33. Sponge Foraging in Dolphins Does Something Unique to Their Genes,
https://www.sciencealert.com/some-dolphins-use-tools-and-it-does-something-unique-to-their-genes 34. Cultural transmission of tool use in bottlenose dolphins,
https://www.researchgate.net/publication/7795611_Cultural_transmission_of_tool_use_in_bottlenose_dolphins 35. Cultural transmission of tool use in bottlenose dolphins - PubMed,
https://pubmed.ncbi.nlm.nih.gov/15947077/ 36. India-centric Hydraulic Civilization - Boloji.com,
https://www.boloji.com/articles/5256/india-centric-hydraulic-civilization 37. Qanat - World History Encyclopedia, https://www.worldhistory.org/qanat/ 38. Qanat: Ancient Water Management in Iran - Groundwater - Scribd,
https://www.scribd.com/document/899184297/Qanat-An-Ancient-Invention-for-Water-Management-in-Iran 39. A simple schematic showing a typical qanat system a Cross-section:,
https://www.researchgate.net/figure/A-simple-schematic-showing-a-typical-qanat-system-a-Cross-section-Using-simple-hand_fig4_369791740 40. Efficiency assessment of some machine learning algorithms for,
https://iwaponline.com/ws/article/25/10/1391/109401/Efficiency-assessment-of-some-machine-learning 41. Sustainable Thermal Energy Storage Systems: A Mathematical,
https://www.researchgate.net/publication/392674872_Sustainable_Thermal_Energy_Storage_Systems_A_Mathematical_Model_of_the_Waru-Waru_Agricultural_Technique_Used_in_Cold_Environments 42. Ancient Agriculture in Harsh Environments - Brewminate,
https://brewminate.com/ancient-agriculture-harsh-environments/ 43. The Thermal Engineering Behind Tiwanaku's Agricultural Success,
https://allthathistory.com/ancient-civilizations/tiwanaku-raised-fields-high-altitude-engineering/4063/ 44. (PDF) Energetic efficiency and political expediency in Titicaca Basin,
https://www.researchgate.net/publication/222200192_Energetic_efficiency_and_political_expediency_in_Titicaca_Basin_raised_field_agriculture