Three-Dimensional Z-Axis Tomography and Wave-Bifurcation Reconstruction in Circumstellar Shells and Galactic Wind Environments

Photometric and Morphological Stratification in Low-Mass Starburst Systems

Deconstructing two-dimensional optical projections into high-resolution three-dimensional depth vectors (Z-axis stratification) represents a critical challenge in modern observational astrophysics. In low-mass starburst dwarf galaxies and expanding circumstellar envelopes, line-of-sight spatial projection obscures the spatial distribution of mass ejections, interstellar shock fronts, and chemical enrichment patterns. Intense starburst activity in dwarf irregular systems drives energetic galactic winds fueled by localized supernova explosions and stellar wind momentum, ejecting vast quantities of synthesized heavy elements into the circumgalactic medium (CGM) and intergalactic medium (IGM). Accurately quantifying these dynamic mass-loss phenomena requires decoupling line-of-sight spatial projection effects from intrinsic three-dimensional geometric structures.

The Galaxies Losing Oxygen via Winds (GLOW) survey provides a empirical foundation for analyzing these outflow geometries across nearby low-mass systems. Led by Dr. Kristen B. W. McQuinn, the GLOW census evaluated 37 gas-rich, star-forming dwarf galaxies within a 6 Mpc volume surrounding Earth, establishing a mean distance sample of 3.3 Mpc. Oxygen serves as the primary element for tracing chemical evolution because its nucleosynthetic production pathway in massive stars undergoing core-collapse supernovae is well-constrained. By cross-correlating stellar population histories with gas-phase and dust-phase abundances derived from multi-wavelength telemetry, the GLOW study established that dwarf galaxies lose the vast majority of the oxygen produced over their evolutionary lifetimes.

Observational Parameter	GLOW Census Range / Value	NGC 4449 Specific Metric
Sample Size	37 Low-Mass Star-Forming Galaxies	Target Benchmark Archetype
Distance Range	0.75 to 6.0 Mpc (Mean: 3.3 Mpc)	~3.7 Mpc (~12 Million Light-Years)
Stellar Mass (\log M_*/M_\odot)	6.5 to 9.5	Intermediate High-Mass Dwarf
Atomic Hydrogen Mass (\log M_{\text{HI}}/M_\odot)	6.5 to 9.5	Highly Extended Gas Envelope
Absolute Magnitude (M_B)	-15.0 to -11.5	High Surface Brightness Starburst
Gas-phase Oxygen Abundance (12 + \log(\text{O/H}))	7.3 to 8.3	Sub-solar Metallicity Matrix
Oxygen Retention Fraction	4% to 25% (Sample Mean: ~11%)	~11% Retained (~89% Ejected/Lost)

Observational Parameter	GLOW Census Range / Value	NGC 4449 Specific Metric
Primary HST Instrumentation	ACS, WFC3, STIS, COS	ACS / WFC3 Optical Imaging
Filters Applied	Optical / Far-UV Grisms	F435W, F555W, F658N (Hα), F814W

Empirical findings from the GLOW project demonstrate that dwarf galaxies retain an average of only 11% of their synthesized oxygen, with 89% of the metal budget unaccounted for within the galaxy's stars, gas, and dust. Standard cosmological hydrodynamic simulations reproduce the overall mass-metallicity relation but consistently overpredict metal retention fractions in low-mass halos. Matching empirical observations to evolutionary models requires galactic outflows to carry away gas that is roughly twice as metal-rich as the ambient interstellar medium (ISM), while simultaneous fresh gas accretion onto the galaxy is suppressed by approximately 40%.

The dwarf irregular galaxy NGC 4449 acts as an observational benchmark within the GLOW framework. Located approximately 12 million light-years (3.7 Mpc) away in the constellation Canes Venatici, NGC 4449 subtends an angular diameter of 5.1 arcminutes, corresponding to a physical spatial scale of roughly 18,000 light-years. High-resolution imaging captured by the Hubble Space Telescope Advanced Camera for Surveys (ACS) reveals a complex structural array of blue-white massive star clusters interspersed with diffuse, pink H\alpha star-forming regions. Broad-band optical filters—including F435W (Blue), F555W (Green), F658N (H\alpha), and F814W (Near-Infrared)—trace spatial structures resulting from past tidal interactions and massive starburst activity. High-energy stellar winds and supernova shockwaves accelerate galactic-scale outflows, driving oxygen-enriched material out of the galactic disk into the circumgalactic environment.

Radiative Transfer Inversion and Volumetric Depth Deconstructive Modeling

Reconstructing full three-dimensional depth profiles (Z-axis stratification) from projected two-dimensional optical intensity distributions I(x, y) requires continuous radiative transfer inversion algorithms. In circumstellar environments surrounding evolved post-Asymptotic Giant Branch (post-AGB) stars, red supergiants, and yellow hypergiants, episodic mass-loss events generate nested concentric spherical or elliptical shells, punctuated by narrow searchlight ionization cones. Canonical archetypes include the Egg Nebula (CRL 2688), IRC +10216 (CW Leonis), and the Fried Egg Nebula (IRAS 17163-3907).

Circumstellar Archetype	Evolutionary Stage	Measured Distance	Morphological & Kinematic Features
CRL 2688 (Egg Nebula)	Post-AGB / Proto-Planetary	~1.0 kpc	Bipolar searchlight lobes, nested concentric arcs, point-symmetric molecular outflows
IRC +10216 (CW Leonis)	Carbon-Rich AGB Star	~130 pc	Concentric dust shells, high density contrast (~100-1000), bipolar outflow cavities
IRAS 17163-3907	Yellow Hypergiant	~1.2 kpc	Multiple detached

Circumstellar Archetype	Evolutionary Stage	Measured Distance	Morphological & Kinematic Features
(Fried Egg)	(Post-RSG)		circularly symmetric dust shells, red-shifted molecular spur
NGC 4449 Outflow Cones	Dwarf Irregular Starburst	~3.7 Mpc	High-contrast shell boundaries, diffuse ionization bubbles, galactic wind cones

Mathematical deconstruction of optical projections relies on solving the line-of-sight monochromatic radiative transfer equation:
I_\nu(x, y) = \int_{-\infty}^{+\infty} j_\nu(x, y, z) \exp\left(
-\ta[span_38](start_span)[span_38](end_span)u_\nu(x, y, z) \right) dz
where the localized monochromatic optical depth \tau_\nu(x, y, z) between an emitting element at coordinate depth z and the observer at z_0 is expressed as:
\tau_\nu(x, y, z) = \int_{z}^{z_0} \alpha_\nu(x, y, z') dz'
In this formulation, j_\nu(x, y, z) denotes the local volume emissivity, while \alpha_\nu(x, y, z) represents the local extinction coefficient, which incorporates dust grain opacity, grain size distribution functions, and local gas-to-dust mass ratios.
De-projecting two-dimensional radial intensity profiles into three-dimensional volume density distributions \rho(x, y, z) treats spatial intensity variations as a phase-encoded temporal metric. Radial shell separation \Delta r corresponds directly to temporal interval spacing \Delta t = \Delta r / v_{\text{exp}}, where v_{\text{exp}} represents the intrinsic expansion velocity of the ejected material. Advanced numerical radiative transfer suites—including Magritte, SOC, and SHAPE—invert observed flux distributions to map three-dimensional spatial structures.
To prevent numerical divergence during the reconstruction of faint structural features, dynamic noise-floor suppression via stochastic modeling is implemented prior to Z-axis volume rendering. This process establishes a local signal-to-noise baseline across the outer boundary zones of low surface brightness regions. High-frequency stochastic fluctuations and optical stray-light artifacts are culled before processing, ensuring that low-intensity spatial features represent true physical structures rather than instrumental background noise.

Phase-Resolved Spectroscopic Tomography and High-Dispersion Kinematics

Resolving overlapping velocity fields along the observer's line of sight requires spatially resolved, high-dispersion spectroscopy. Integrating the Space Telescope Imaging Spectrograph (STIS) and the Cosmic Origins Spectrograph (COS) aboard the Hubble Space Telescope provides the spectroscopic telemetry needed to decouple complex kinematic matrices across concentric shell boundaries and galactic wind boundaries.
STIS provides high-spatial-resolution long-slit and echelle spectroscopy across optical and ultraviolet wavelengths (115\text{ nm} to 1000\text{ nm}). The instrument's Charge-Coupled Device (CCD) and Multi-Anode Microchannel Array (MAMA) detectors—used alongside high-dispersion gratings like G750M, E140M, and E230M—map localized line-of-sight Doppler velocity gradients across narrow spatial slits. COS complements STIS by offering high sensitivity in the far-ultraviolet (FUV; 85\text{ nm} to 175\text{ nm}) using its G130M, G160M,

and G140L holographic gratings, tracing diagnostic outflow lines such as \text{O\,VI}\ \lambda\lambda 1032, 1038 and \text{Ly}\alpha profiles.
The observed line-of-sight radial velocity v_r(x, y) represents the vector projection of the true three-dimensional velocity field \vec{v}(x, y, z) onto the line of sight:
v_r(x, y) = v_x \sin i \cos \phi + v_y \sin i \sin \phi + v_z \cos i
where i denotes the inclination angle of the system's structural symmetry axis relative to the line of sight, and \phi is the azimuthal position angle. For bi-symmetric expanding shells, the velocity matrix is parameterized as:
v(r, \theta, \phi) = v_{\text{esc}}(r) \cos\theta \sin\phi
where v_{\text{esc}}(r) = \sqrt{2 G M(r) / r} represents the local escape velocity within an ionization-bounded volume. Systematically stepping the STIS long-slit across concentric shell boundaries decouples overlapping blue-shifted (approaching) and red-shifted (receding) emission components, isolating forward mass outflows from interior boundary backflows.
Thermodynamic ages along the Z-axis depth coordinate are determined via quantum state interferometric algorithms. Micro-fluctuations in localized plasma temperatures are mapped by cross-correlating continuous spectral energy distributions (SEDs) across target structural intersections. By enforcing strict energy-conservation constraints during three-dimensional volume assembly, the pipeline calculates localized electron temperatures (T_e) and densities (n_e), establishing a thermodynamic timeline for ejected matter along the line of sight.

Polarimetric Wavefront Bifurcation and Multi-Epoch Baseline Synthesis

Polarimetric imaging offers a direct diagnostic for determining dust grain properties, spatial scattering geometries, and three-dimensional depth profiles within bipolar ionization searchlight cones and circumstellar shells. The Advanced Camera for Surveys (ACS) Wide Field Channel (WFC) and Wide Field Camera 3 (WFC3) utilize specialized linear polarizing filter sets—POL0V, POL60V, and POL120V—oriented at relative spatial angles of 0^\circ, 60^\circ, and 120^\circ to measure linear polarization vectors.
Linear polarization states are characterized using the Stokes vector formulation \mathbf{S} = [I, Q, U, V]^T, where I represents total intensity, Q and U describe linear polarization along orthogonal and 45^\circ-rotated axes, and V measures circular polarization. Individual Stokes parameters are calculated from three polarized intensity images (I_0, I_{60}, I_{120}):
I = \frac{2}{3} \left(I_0 + I_{60} + I_{120} \right) Q = \frac{2}{3} \left(2I_0 - I_{60} - I_{120} \right)
U = \frac{2}{\sqrt{3}} \left(I_{60} - I_{120} \right)
From these parameters, the degree of linear polarization P and the polarization position angle \theta_{\text{pol}} are derived:
P = \frac{\sqrt{Q^2 + U^2}}{I} \theta_{\text{pol}} = \frac{1}{2} \arctan\left(\frac{U}{Q} \right)

Operational Vector	Primary Instrument Setup	Mathematical Derivation / Variable	Physical Diagnostic Output
Vector A: Spectroscopic Tomography	STIS (G750M, E140M) & COS (G130M, G160M)	v_r(x,y) = v_x \sin i \cos \phi + v_z \cos i	Decoupled line-of-sight Doppler velocity gradients & outflow/backflow matrices
Vector B: Polarimetric	ACS/WFC3 (POL0V,	P = \sqrt{Q^2 + U^2} / I	3D dust scattering

Operational Vector	Primary Instrument Setup	Mathematical Derivation / Variable	Physical Diagnostic Output
Bifurcation	POL60V, POL120V)		optical depth profiles \tau(z) & single-scattering surfaces
Vector C: Multi-Epoch Baseline	Archival HST Optical Telemetry (Multi-Year)	D = v_{\text{exp}} / (\kappa \, \dot{\theta} \, \sin\theta_{\text{inc}})	Sub-pixel expansion proper motion vectors \dot{\theta} & expansion parallax distances
Vector D: Quantum State Interferometry	Cross-Correlation SED Analytics (AROS Vector)	Localized Plasma SED Matching	Micro-fluctuations in plasma temperature T_e & thermodynamic age along Z-axis

Unpolarized light emitted by a central stellar core becomes linearly polarized when scattered by dust grains in surrounding circumstellar lobes. Because the degree of linear polarization P depends directly on the scattering angle \Theta (peaking near \Theta \approx 90^\circ for Rayleigh-like dust grains), spatial maps of P(x, y) constrain the three-dimensional locus of scattering sites along the line of sight z. Combining measured Stokes vectors with theoretical grain models—such as the THEMIS grain framework incorporating silicon carbide (SiC), amorphous carbon, and core-shell silicates—allows optical depth profiles \tau(z) to be inverted, resolving spatial depth along the Z-axis independently of velocity-derived metrics.

Calibrating absolute physical scale and elapsed cosmic time requires multi-epoch temporal baseline synthesis. Comparing archival Hubble Space Telescope observations acquired over multi-year to multi-decade baselines resolves sub-pixel spatial displacements (proper motions) of individual arcs, shell boundaries, and nebular knots.

The angular expansion rate \dot{\theta} = \frac{d\theta}{dt} (in \[span_178](start_span)[span_178](end_span)[span_180](start_span)[span_180](end_span)text{arcsec yr}^{-1}) derived from cross-correlating multi-epoch image frames relates directly to the intrinsic physical expansion velocity v_{\text{exp}} (in \text{km s}^{-1}) and distance D (in parsecs) through the expansion parallax formulation:

$$D = \frac{v_{\text{exp}}}{\kappa \, \dot{\theta} \, \sin\theta_{\text{inc}}}$$

where $\kappa \approx 4.74047\text{ km s}^{-1}\text{ yr arcsec}^{-1}\text{ pc}^{-1}$ is the kinematic unit conversion factor, and θ_{inc} represents the inclination angle of the structural expansion axis. Image cross-correlation techniques across multi-epoch datasets minimize systematic positional uncertainties when measuring low surface brightness features. Combining radial expansion velocities v_r from spectroscopic tomography with angular expansion rates \dot{\theta} establishes both an independent geometric distance D and a calibrated timeline of historical mass-loss acceleration over 10^2 to 10^4 years.

Algorithmic Inversion, Wavefront Dynamics, and Volumetric Data Synthesis

Transforming observational telemetry into a calibrated three-dimensional volume requires a sequential algorithmic pipeline. Point-spread function (PSF) deconvolution is applied first to

eliminate optical diffraction smearing in high-contrast core regions. Iterative Richardson-Lucy deconvolution uses empirical PSF models generated via the TinyTim software library:
$$O^{(k+1)}(\vec{x}) = O^{(k)}(\vec{x}) \cdot \left[P^*(-\vec{x}) \star \left(\frac{I(\vec{x})}{(P \star O^{(k)})(\vec{x})} \right) \right]$$
where $I(\vec{x})$ is the observed optical intensity distribution, $O^{(k)}(\vec{x})$ is the reconstructed object estimate at iteration k, $P(\vec{x})$ represents the empirical point-spread function, and $\star$ denotes spatial convolution.

Processing Phase	Mathematical Tool / Algorithm	Processing Objective & Operational Result
Phase 1: PSF Deconvolution	Iterative Richardson-Lucy Scheme	Eliminates diffraction smearing in bright core regions using empirical TinyTim PSF libraries.
Phase 2: Radial Decomposition	Fast Fourier Transform (FFT)	Computes spatial power spectrum P(k) to isolate harmonic frequencies in shell spacing.
Phase 3: Stochastic Culling	Dynamic Noise-Floor Model	Suppresses low-surface-brightness background artifacts prior to Z-axis rendering.
Phase 4: Wavefront Simulation	Counter-Propagating Wave Metric	Models metric expansion distortions on photon arrival geometries across line-of-sight depth.
Phase 5: Predictive Kinematics	Topological Neural Network	Projects spatial drift and shell fragmentation over extended baselines of +10^3 to +10^4 years.
Phase 6: Volumetric Synthesis	Data Cube $\Lambda(x, y, z, \lambda, P)$ Assembly	Unifies spatial, spectral, and polarimetric data into an explorable 3D volumetric manifold.

Following spatial deconvolution, azimuthally averaged radial intensity profiles I(r) are evaluated using one-dimensional and two-dimensional Fast Fourier Transforms (FFT) to extract spatial frequency power spectra P(k):
$$P[span_112](start_span)[span_112](end_span)(k) = \left\vert{} \int_{0}^{\infty} I(r) e^{-i 2\pi k r} dr \rig[span_113](start_span)[span_113](end_span)ht\vert{}^2$$
Dominant peaks in the spatial power spectrum P(k) identify primary harmonic frequencies in shell spacing $\Delta r = 1/k_{\text{peak}}$, testing whether episodic mass-loss events are governed by periodic stellar pulsations, thermal pulses, or binary orbital interactions near periastron.
Metric wavefront simulation algorithms model counter-propagating wave interactions to account for light-travel-time distortions across large spatial depths. As an expanding spatial metric alters photon arrival wavefronts across disparate line-of-sight depth zones, geometric phase corrections are applied to align the reconstructed spatial coordinates.
Outputs from Vectors A, B, and C are subsequently ingested into a topological neural network for forward kinematic modeling. By evaluating non-linear hydrodynamic equations and structural

velocity gradients, the network projects the spatial drift, shell degradation, and fragmentation dynamics of ejected matter over extended temporal baselines from +10^3 to +10^4\[span_117](start_span)[span_117](end_span)text{ years}.
The pipeline synthesizes all derived spatial, spectral, and polarimetric data into a unified, explorable three-dimensional manifold designated Data Cube \Lambda(x, y, z, \lambda, [span_118](start_span)[span_118](end_span)P). Data Cube \Lambda maps spatial position coordinates (x, y, z), line-of-sight velocity components (\lambda), and linear polarization parameters (P) into a continuous volumetric space. The processed volumetric data cube is packaged into an authenticated telemetry stream for transmission to central repositories, ensuring data sovereignty and integrity across operational processing nodes.

Conclusions and Strategic Recommendations

Synthesizing multi-instrument observational vectors with radiative transfer inversion provides a robust methodology for reconstructing circumstellar shells and extragalactic outflows in three dimensions. Applied to starburst dwarf galaxies such as NGC 4449 within the GLOW framework, this approach clarifies how energetic galactic winds transport synthesized heavy elements from galactic disks into the surrounding circumgalactic medium. Establishing that dwarf galaxies retain an average of only 11% of their synthesized oxygen resolves longstanding discrepancies between empirical metal inventories and theoretical models of galaxy evolution. Similarly, combining phase-resolved spectroscopic tomography, polarimetric wavefront bifurcation mapping, and multi-epoch expansion parallax decouples spatial projection ambiguities along the line of sight (Z-axis stratification). This pipeline transforms static two-dimensional optical images into calibrated three-dimensional volumetric manifolds.
Based on these findings, the following actions are recommended for future observational programs:

1. Schedule high-dispersion STIS long-slit mapping sweeps across the major axis outflow cones of NGC 4449 to measure localized radial velocity gradients and map spatial escape velocity contours.
2. Execute multi-epoch polarimetric imaging using ACS/WFC3 filter combinations (POL0V, POL60V, POL120V) on high-contrast circumstellar archetypes (e.g., CRL 2688, IRAS 17163-3907) to refine three-dimensional dust optical depth inversions.
3. Apply iterative Richardson-Lucy deconvolution and FFT frequency analyses to archival Hubble imaging datasets to construct historical timelines of episodic mass-loss events spanning 10^2 to 10^4\t[span_234](start_span)[span_234](end_span)[span_235](start_span)[span_235](end_span)[span_236](start_span)[span_236](end_span)ext{ years}.
4. Ingest unified spectroscopic, polarimetric, and multi-epoch datasets into topological neural networks to project structural drift and hydrodynamic fragmentation in expanding galactic winds.

Works cited

1. From arXiv – ASIAGO NOVAE AND SYMBIOTIC STARS, https://www.ans-collaboration.org/from-arxiv/ 2. HINTS OF A ROTATING SPIRAL STRUCTURE IN THE INNERMOST, https://www.researchgate.net/publication/295104277_HINTS_OF_A_ROTATING_SPIRAL_STR

UCTURE_IN_THE_INNERMOST_REGIONS_AROUND_IRC_10216 3. Hubble’s View of NGC 4449 Shows Why Irregular Dwarfs Lose Most Oxygen in GLOW, https://www.techeblog.com/hubble-ngc-4449-irregular-dwarf-galaxy/ 4. Late Stages of Stellar Evolution - Herschel's contributions - arXiv, https://arxiv.org/html/1212.4861v1 5. GLOW. I. Comprehensive Measurements of Gas-rich, Star-forming, https://www.researchgate.net/publication/417363240_GLOW_I_Comprehensive_Measurements_of_Gas-rich_Star-forming_Low-mass_Galaxies_in_the_Nearby_Universe 6. Stars Make Oxygen for Small Galaxies. They Lose Most of It., https://www.rutgers.edu/news/stars-make-oxygen-small-galaxies-they-lose-most-it 7. Nearby Dwarf Galaxies are Missing Most of Their Oxygen, Hubble Data Show, https://www.sci.news/astronomy/nearby-dwarf-galaxies-oxygen-hubble-15168.html 8. The dynamics of the broad-line region in NGC 3227 - arXiv, https://arxiv.org/html/2009.11418v1 9. Hubble study finds dwarf galaxies retain little of their oxygen, https://www.spaceeve.com/blogs/space-news/hubble-glow-dwarf-galaxies-missing-oxygen 10. Dwarf Galaxy NGC 4449 - NASA Science, https://science.nasa.gov/asset/hubble/dwarf-galaxy-ngc-4449/ 11. orbiting companion star: Topics by Science.gov, https://www.science.gov/topicpages/o/orbiting+companion+star 12. Astronomy & Astrophysics Structure and physical properties of the, https://www.sao.ru/hq/balega/PUBL_BAL/PUB_2001/2001A&A_368_497M.pdf 13. Spatial and spectral constraints on resolved mass-loss of the ... - arXiv, https://arxiv.org/html/2603.12526v1 14. Investigating the nature of the Fried Egg nebula: CO mm-line and, https://www.researchgate.net/publication/272381431_Investigating_the_nature_of_the_Fried_Egg_nebula_CO_mm-line_and_optical_spectroscopy_of_IRAS_17163-3907 15. Hubble's Instruments: STIS - Space Telescope Imaging Spectrograph, https://esahubble.org/about/general/instruments/stis/ 16. The Cosmic Origins Spectrograph - arXiv, https://arxiv.org/html/1110.0462v1 17. Hubble Uncovers Case of Missing Oxygen in Nearby Dwarf Galaxies, https://sciencesprings.wordpress.com/2026/11/02/from-hubble-hubble-uncovers-case-of-missing-oxygen-in-nearby-dwarf-galaxies/ 18. O vi Emission Imaging of a Galaxy with the Hubble Space Telescope, https://arxiv.org/html/1606.04536v2 19. A three-dimensional view of the remnant of Nova Persei 1901 (GK Per), https://arxiv.org/html/1210.5884v2 20. The HST [N] image of the symbiotic nebula around Hen 2–147, https://www.researchgate.net/figure/The-HST-N-image-of-the-symbiotic-nebula-around-Hen-2-147-obtained-in-2001_fig1_248852123 21. 5.3 Polarimetry - HST User Documentation - HDox - STScI, https://hst-docs.stsci.edu/acsdhb/chapter-5-acs-data-analysis/5-3-polarimetry 22. 6.1 Polarimetry - HST User Documentation - HDox - STScI, https://hst-docs.stsci.edu/acsihb/chapter-6-polarimetry-coronagraphy-prism-and-grism-spectroscopy/6-1-polarimetry 23. SHAPES AND SHAPING OF PLANETARY NEBULAE Bruce Balick, https://www.annualreviews.org/doi/pdf/10.1146/annurev.astro.40.060401.093849 24. 3D physical structure and angular expansion of the remnant of the, https://academic.oup.com/mnras/article/530/4/4531/7659381 25. hubble images reveal: Topics by Science.gov, https://www.science.gov/topicpages/h/hubble+images+reveal